

Case SCARBAN combined with ALHYDRAN: Dog bite

1. Case study description

patient age	10 years
gender	boy
cause	dog bite
why scar treatment	<ul style="list-style-type: none"> - no expectation of scar improvement - increase in hypertrophy - poor mobility of the finger (dig 3 left) - persistent itching and redness
start scar treatment	3 months after bite
therapy	6 months of intensive scar treatment: <ul style="list-style-type: none"> - SCARBAN® Finger (including pressure stocking) at night - ALHYDRAN cream (2 to 3 times daily) during the day

2. Evolution of the scar



Open wound

After 3 months:
start of scar treatmentScar after 3 months of
scar treatmentScar after 6 months of
scar treatment

3. Conclusions medical team

At the start of treatment, the scar looked bad: reactive and no signs of improvement.
Within 3 months of scar treatment improvements are already visible.

ALHYDRAN by day:

- direct cooling effect
- during the treatment itching and redness reduced and skin became more supple

SCARBAN® Finger at night:

- scar became more relaxed and smoother
- improved mobility of the hand
- less itching, pain and redness

4. Patient experience

In order not to restrict the 10-year-old boy at school and while playing, a combined treatment (ALHYDRAN medical cream + SCARBAN® silicone scar treatment) was prescribed.

Author: Ms L. Rolfes-van Os, Hand and Occupational Therapist, Revalidatiecentrum Saxenburg, Hardenberg, the Netherlands